

*On Satellite Constellation Selection*

DISTRIBUTION STATEMENT A  
Approved for public release;  
Distribution Unlimited

19980513 332

**DTIC QUALITY INSPECTED 4**

PLEASE RETURN TO:

BMD TECHNICAL INFORMATION CENTER  
BALLISTIC MISSILE DEFENSE ORGANIZATION  
7100 DEFENSE PENTAGON  
WASHINGTON D.C. 20301-7100

Los Alamos

U3361

*Los Alamos National Laboratory is operated by the University of California for the United States Department of Energy under contract W-7405-ENG-36.*

*Prepared by Bo West, P Division*

*An Affirmative Action/Equal Opportunity Employer*

*This report was prepared as an account of work sponsored by an agency of the United States Government. Neither The Regents of the University of California, the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by The Regents of the University of California, the United States Government, or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of The Regents of the University of California, the United States Government, or any agency thereof.*

Accession Number: 3361

Publication Date: May 01, 1991

Title: On Satellite Constellation Selection

Personal Author: Canavan, G.H.

Corporate Author Or Publisher: Los Alamos National Laboratory, Los Alamos, NM 87545 Report Number: LA-12059-MS

Report Prepared for: US Department of Energy

Descriptors, Keywords: Satellite Distribution Constellation Optimization Inclination Orbit

Pages: 00007

Cataloged Date: Jan 31, 1992

Contract Number: W-7405-ENG-36

Document Type: HC

Number of Copies In Library: 000001

Record ID: 23191

*On Satellite Constellation Selection*

*Gregory H. Canavan*

PLEASE RETURN TO:  
SDI TECHNICAL INFORMATION CENTER

## CONTENTS

ABSTRACT	1
I. INTRODUCTION	1
II. SATELLITE DISTRIBUTION	2
III. CONSTELLATION OPTIMIZATION	3
A. Single Inclination	3
B. Multiple Inclinations	3
IV. SUMMARY AND CONCLUSIONS	4
ACKNOWLEDGMENT	5
REFERENCES	6

# ON SATELLITE CONSTELLATION SELECTION

by

Gregory H. Canavan

## ABSTRACT

Analytical estimates can be used to produce and discuss optimal constellations. They are in close agreement with phase-space estimates and exact solutions. They suggest that distributions of inclined orbits could reduce satellite numbers by factors of 2-3 while improving uniformity.

---

## I. INTRODUCTION

In evaluating satellite constellations, an essential element is their ability to produce uniform distributions of satellites over the latitudes of interest, although practical considerations can be overriding. This report discusses some of the results available for evaluating constellations such as communications networks that require some degree of uniformity for practicality. Both computer simulations and analytic solutions are available. Numerical simulations are the most accurate, but analytic estimates give more insight into the sensitivity of results to parameter and missile distributions.

## II. SATELLITE DISTRIBUTION

Satellite orbits can be taken to be circular. Their time and longitudinally averaged distributions are used below. A satellite whose orbit is inclined an angle  $i \leq \pi/2$  from the north pole remains in a band  $-i \leq \theta \leq i$  of latitude. If  $\theta$  is the satellite's latitude and  $\epsilon$  its azimuthal angle around its pole, from the spherical law of cosines

$$\sin\theta = \sin i \cdot \cos\epsilon, \quad (1)$$

where  $\epsilon = 0$  for  $\theta = i$ . In traversing a latitude band  $d\theta$  the satellite moves through an angle around its pole of<sup>1</sup>

$$d\epsilon = \cos\theta \cdot d\theta / (\sin^2 i - \sin^2 \theta)^{1/2}, \quad (2)$$

traversing a fraction  $d\epsilon/2\pi$  of its orbit in the process. This latitude band contains a fraction  $\cos\theta \cdot d\theta/2$  of the earth's area. If the orbit contains 1 satellite, the satellite density is

$$n_+(\theta) = (d\epsilon/2\pi) / 2\pi \cos\theta \cdot d\theta = 1/4\pi^2 (\sin^2 i - \sin^2 \theta)^{1/2}, \quad (3)$$

where distance is measured in multiples of the earth's radius  $R_e$  and  $n_+$  is the density of satellites ascending in latitude. By symmetry, at any latitude there is an equal density,  $n_-$ , of satellites descending, and the total density is their sum<sup>2</sup>

$$n(\theta) = n_+ + n_- = 1/2\pi^2 (\sin^2 i - \sin^2 \theta)^{1/2}. \quad (4)$$

$n(\theta)$  is fairly flat, diverging integrably at  $\theta = i$ , but  $n(\theta)$  is much larger at high than low latitudes.

Figure 1 shows the satellite densities as functions of latitude for constellations with single inclinations of 60, 70, 80, and 90 degrees. For polar orbits the satellites pile up over the pole. For smaller inclinations the variations are less, although the nonuniformities still approach factors of two at the northern limits of coverage.

As an aside, for single inclinations, satellites are moving in just two directions, ascending or descending, at any latitude. Since  $n_+ = n_-$ , their component densities are  $n_+ = n_- = n(\theta)/2$  at each latitude. The angle between ascending orbits and the local east is

$$\Omega_+(\theta, i) = \sin^{-1}(d\theta/d\epsilon) = \sin^{-1}[(\sin^2 i - \sin^2 \theta)^{1/2} / \cos\theta]. \quad (5)$$

For descending satellites the angle is  $\Omega_- = -\Omega_+$ . They cross at an angle of roughly  $2\Omega_+$ . At the orbit's maximum latitude  $\Omega_+ = \Omega_-$

= 0; both ascending and descending orbits are tangent to the maximum latitude  $i$  there. At the equator,  $\Omega_- = -i$ , so the ascending satellites cross circles of constant latitude at monotonically decreasing angles.

### III. CONSTELLATION OPTIMIZATION

This section uses the satellite distributions above to derive effective distributions of inclinations. A uniform distribution of satellites,  $n_u = 1/4\pi$ , for which the fraction of satellites between latitudes  $0 \leq \theta_1 \leq \theta \leq \theta_2$  is

$$f_u = \int_{\theta_1}^{\theta_2} d\theta \cdot 2\pi \cos\theta \cdot (1/4\pi) = (\sin\theta_2 - \sin\theta_1)/2, \quad (6)$$

which is the basis for comparison below.

#### A. Single Inclination

For a constellation with a single inclination  $i$ , Eq. (4) can be integrated to obtain the fraction of the satellites in any given latitude band, which is

$$f_i = \int d\theta \cdot 2\pi \cos\theta / 2\pi^2 (\sin^2 i - \sin^2 \theta)^{1/2}. \quad (7)$$

When the limits on the integral are  $\theta_1$  and  $\theta_2$ , Eq. (7) gives

$$f_i = [\sin^{-1}(\sin\theta_2/\sin i) - \sin^{-1}(\sin\theta_1/\sin i)]/\pi. \quad (8)$$

The average constellation concentration possible over  $\theta_1$  to  $\theta_2$  is  $z \equiv f_i/f_u$ , whose maximum occurs at  $i = \theta_2$ ,<sup>3</sup> where it is

$$z = f_i/f_u = [1 - (2/\pi) \sin^{-1}(\sin\theta_1/\sin\theta_2)] / (\sin\theta_2 - \sin\theta_1). \quad (9)$$

Both  $n$  and  $z$  vary rapidly over high-latitude bands, which makes estimates sensitive to the averaging and specific values used.

#### B. Multiple Inclinations

The maximization of constellation concentration has been extended to multiple inclinations, which improve uniformity. The starting point is the generalization of Eq. (4) to multiple inclinations, which is

$$n(\theta) = \int di \cdot N(i) / 2\pi^2 (\sin^2 i - \sin^2 \theta)^{1/2}, \quad (10)$$

where the integral is over the inclinations for which satellites can reach latitude  $\theta$ . For a launch area within  $\theta_1 \leq \theta \leq \theta_2$ , the objective is to choose  $N(i)$  such that  $n(\theta)$  is constant in that interval. The solution to this Abel equation is<sup>4</sup>



$$N(i) = \cos i \cdot \sin i / \sqrt{(\cos^2 i - \cos^2 \theta_2)} \quad (11)$$

for  $\theta_1 \leq \theta \leq \theta_2$  and 0 elsewhere. For a uniform distribution over  $\theta_1 = 0 \leq \theta \leq \theta_2 = \pi/2$  Eq. (11) reduces to  $N(i) = \sin i$ , for which Eq. (10) gives  $n(\theta) = 1/4\pi$  for all  $\theta$  and  $\Sigma di \cdot N(i) = 1$ , which is the geometric result for a random distribution of satellites.

The integral of  $N(i)$  over  $\theta_1 \leq i \leq \theta_2$  gives the fraction of the satellites in that interval. Dividing the geometric fraction from the uniform satellite distribution to that produced by the inclination distribution of Eq. (11) gives

$$z_w = 1 / \sqrt{(\cos^2 \theta_1 - \cos^2 \theta_2)}. \quad (12)$$

This concentration factor is  $z_w \approx 2.5$  for  $n$  uniform between  $\theta_1 = 50^\circ$  and  $\theta_2 = 60^\circ$ , which is  $\approx 20\%$  less than the  $z \approx 3$  from the non-uniform single inclination of Eq. (9).

The reduction in the number of satellites by using a distribution of inclinations rather than a single one is given by the ratio  $f_i/f_w$ . For coverage down to the equator  $f_w \rightarrow 1/2$ ,  $f_i \rightarrow \sin^{-1}(\sin \theta_2 / \sin i) / \pi$ . For  $i = \theta_2 = \pi/2$ , coverage to the pole, the ratio  $f_i/f_w$  is unity. For polar orbits  $f_i \rightarrow \theta_2/\pi$ . Thus, for  $\theta_2 = \pi/4$ , i.e., coverage of the U.S. and lower Europe down to the equator,  $f_i \rightarrow 1/4$ , and the gain for a distribution of inclinations is a factor of 2. Other combinations are shown in Fig. 2. For coverage up to  $60^\circ$ , to include the Soviet Union, polar orbits would pay about a factor of 2 in satellites over a distribution up to  $60^\circ$ . Thus, distributed constellations are significantly smaller as well as more uniform.

#### IV. SUMMARY AND CONCLUSIONS

Analytical estimates can be used to produce and discuss optimal constellation estimates for uniform coverage. Those estimates are in reasonable agreement with phase space estimates and exact solutions. They suggest that using distributions of inclined orbits could reduce the number of satellites by factors of 2-3, while improving uniformity.

#### ACKNOWLEDGMENT

The author would like to acknowledge stimulating discussions of constellation scaling with Dr. Albert Petscheck and of constellation selection with Dr. Raymond Leopold of Motorola.

## REFERENCES

1. A. Petschek, Satellite Density Calculation, Los Alamos National Laboratory memo, 10 May 1985.
2. R. Garwin, "How Many Orbiting Lasers for Boost-Phase Intercept?" Nature, 315, 23 May 1985, pp. 286-90.
3. A. Petschek, op cit.
4. G. Wing, Los Alamos National Laboratory memo C-3-1097D, 13 June, 1985.

Fig. 1 Satellite density vs inclination





Fig. 2 Uniform vs inclined constel.



This report has been reproduced directly from the best available copy.

It is available to DOE and DOE contractors from the Office of Scientific and Technical Information, P.O. Box 62, Oak Ridge, TN 37831. Prices are available from (615) 576-8401, FTS 626-8401.

It is available to the public from the National Technical Information Service, U.S. Department of Commerce, 5285 Port Royal Rd., Springfield, VA 22161.