
# LEGIBILITY NOTICE

A major purpose of the Technical Information Center is to provide the broadest dissemination possible of information contained in DOE's Research and Development Reports to business, industry, the academic community, and federal, state and local governments.

Although a small portion of this report is not reproducible, it is being made available to expedite the availability of information on the research discussed herein.

1



Los Alamos National Laboratory is operated by the University of California for the United States Department of Energy under contract W-7405-ENG-36

LA-UR--90-2212

DE90 015104

TITLE: FISSION-PRODUCT CHAIN YIELDS AND DELAYED NEUTRONS: ANS 5.2 AND ANS 5.8

AUTHOR(S): T. R. England, T-2

B. F. Rider, T-2 Consultant .

M. C. Brady, Oak Ridge Nat. Lab.

SUBMITTED TO:

The American Nuclear Society for presentation at the 1990 Winter Meeting, Washington D.C., Nov. 11-15, 1990.

#### DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufactures, or adieswise loss not necessarily constitute or imply its enforsement, recomme idation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof

Cylandepranse of this article, the publisher recognites that the U.S. Government retains a none-clusive, injustry free ficeuse to publish or reproduce may be used force of the contribution or to allow others to do so, for U.S. Government purposes

than process that the formatory requests that the publisher objudy this article as work performed number the anspices of the U.S. Department of coercy



# FISSION-PRODUCT CHAIN YIELDS AND DELAYED NEUTRONS: ANS 5.2 AND ANS 5.8

T. R. England Los Alamos National Laboratory

B. F. Rider General Electric Co. Retiree, Los Alamos Nat. Lab. Consultant

M. C. Brady
Oak Ridge National Laboratory

### YIELDS (ANS 5.2)

Chain yields are the addition of the direct values along constant (Z+N) paths (see Fig. 1). The addition must also account for decay branching, especially for delayed neutrons that couple the mass chains. The result is the familiar double-humped plot of yield per fission vs A (also evident in Fig. 1). The lines of stable nuclides and most probable yield are shown in the (Z,N) plane. Some modern measurements provide direct yields; others provide cumulative values for long-lived or stable products. Yield evaluations must account for each type of measurement and the degree of decay coupling, beginning with direct yields.

For some fissioning nuclides at thermal or fast neutron incident energies, the amount of data is enormous; and for other nuclide-energy combinations, the data must be developed from systematics. 1,2,3 Many applications of chain yields can use the same systematics to estimate independent values. Reference 3 will include all measurements (1939-1990) and recommended yields (that include measurements as well) for sixty yield sets, each containing approximately 1200 fission products. However, the document will contain over 9000 pages: its equivalent in ENDF/B-VI will contain over 250 000 card images. Many users will prefer to refer to the ANS 5.2 chain yield standard, depending upon each application.

The ANS 5.2 standard is currently in rough draft form for comment from Working Group members. It includes 107 mass chain yields and uncertainties for <sup>233,235,238</sup>U, <sup>239,240,241</sup>Pu, <sup>232</sup>Th, and <sup>252</sup>Cf. Only fast fission yields are included for <sup>232</sup>Th, <sup>238</sup>U, and <sup>240</sup>Pu, and spontaneous yields for <sup>252</sup>Cf. The final chain yields in it will be developed from (but not included in) the individual nuclide yields and branchings in ENDF/B-VI. A relatively small ANS document can be generated due to the existence of ENDF/B-VI and inclusion of measurement, and references (through April 1990), as seen in Ref. 3.

Some applications of this standard and problems, e.g., with energy dependence, will be discussed.

### **DELAYED NEUTRONS (ANS 5.8)**

The first meeting of this Working Group occurred on 10 June 1990. A related international working group was formed by NEACRP/NEANDC in April 1990 to provide delayed neutron benchmarks. Several individuals have joint membership with the ANS 5.8 group and a good level of cooperation is expected. The international group is concerned primarily with  $\beta_{eff}$ ,  $\vec{v}_d$ , six-group data, and the energy dependence of these, but this will almost certainly expand to include precursor data.

We decided to keep precursor data (Pn and spectra),  $\bar{v}_d(E)$ , and six-group data (spectra, abundance and halflives) within the scope of the ANS 5.8 standard, unless it later proves impossible to achieve an acceptable consistency with benchmarks when using precursor data.

The reader should be aware that, while ANS 5.8 is very new, the effort to improve delayed neutron precursor data has been progressing nearly continuously since the mid-70's, with a particularly large effort expending during 1984-87. Such data are necessary for ENDF decay files as well as yield files. However, except for Pn data in ENDF/B-V, individual precursor data were not included in this version. We found the aggregate six-group spectra to be inadequate and decided to improve these spectra with precursor calculations. F. Mann et al.<sup>6</sup> evaluated measured Pn values and T. England et al.<sup>7</sup> updated spectra with available measurements. Later (> 1984) we decided that the measured precursor spectra were still inadequate in energy range and in the number of precursors having measured spectra. Simple models were used to expand the spectra where measurement exist and to provide spectra for precursors having no measurements.<sup>8</sup> Pn values were estimated from systematics for unmeasured values. A consistent set of six-group abundances, halflives, and spectra were developed from the resulting 271 precursor data base. Brady et al.<sup>8</sup> provide a particularly good description of this work, including comparisons with very recent measurements.<sup>9</sup>

The six-group data will be incorporated into ENDF/B-VI, thereby extending the number of fissioning systems from 7 to 28 and expanding the spectra from  $\sim 79 \text{ keV} - 1.2 \text{ MeV}$  to 0 - 3 MeV. Individual precursor spectra and Pn values for 271 nuclides will also be incorporated into the ENDF/-VI decay files. Some current Pn values may be altered as

a result of recent Studsvik measurements (reported in Ref. 10).  $\overline{v}_d$  data remain as the values determined by  $Cox^4$  and Tuttle,<sup>5</sup> or those calculated where  $\overline{v}_d$  had not been measured, with a few exceptions.<sup>8</sup>

We were not aware of the European concern with inconsistent  $\beta_{eff}$  measurements until the ENDF/B-VI data were completed. The primary reason for this concern is based on  $^{238}$ U fast fission. Unfortunately, this is also the most inconsistent case of precursor calculations and measurements. Perhaps a reason and solution of the problem will be determined by the time of this meeting.

In summary, ANS 5.2 is near completion and ANS 5.8 is starting from existing, extensive work, with the cooperation of the international community involved.

## REFERENCES:

- 1. T. R. England and J. Blachot, "Status of Fission Yield Data," Proc. Int. Conf. on Nuclear Data for Sci. and Tech., May 30-June 3, 1988, Mito, Japan.
- 2. A. C. Wahl, "Nuclear-Charge and Mass Distribution from Fission," Proc. Conf."50 years with Nuclear Fission," Washington, D.C., April 25-28, 1989, J. W. Behrens and A. D. Carlson, Eds.
- 3. T. R. England and B. F. Rider, "Evaluation and Compilation of Fission-Product Yields, 1989," ENDF-349 [draft completed; report to be published by Los Alamos National Laboratory, 1990].
- 4. S. A. Cox, "Delayed Neutron Data--Review and Evaluation," Argonne National Laboratory report ANL/NDM-5 (April 1974). [The data were updated for ENDF/B-V by R. E. Kaiser and S. G. Carpenter, Argonne National Laboratory-West (unpublished).]
- 5. R. J. Tuttle, "Review of Delayed Neutron Yields in Nuclear Fission," Proc. Consultants' Mtg. Delayed Neutron Properties, Vienna, Austria, March 26-30, 1979, INDC-NDS-107/G ± Special, p. 29, Int. Atomic Energy Agency (1979).
- 6. F. M. Mann, M. Schreiber, R. E. Schenter, and T. R. England, Nucl. Sci. Eng. 87, 418 (1984). [See also update by F. M. Mann, Proc. Specialists' Mtg. Delayed Neutrons, Birmingham, England, September 15-19, 1986, ISBN 07044-09267, D. R. Weaver, Ed., Univ. of Birmingham.]
- 7. T. R. England, W. B. Wilson, R. E. Schenter, and F. M. Mann, Nucl. Sci. Eng. 85, 139 (1983).
- 8. M. C. Brady and T. R. England, Nucl. Sci. Eng. <u>J03 (2)</u>, 109 (1989). [See also thesis by M. C. Brady, "Evaluation and Application of Delqayed Neutron Precursor Data," Los Alamos National Laboratory Thesis Series report LA-1153-T (April 1989).]

- 9. G. Couchell, P. Bennet, E. Jacobs, D. Pullen, R. Tanczyn, M. Haghighi, and Q. Sharfuudin, "Composite Delayed Neutron Energy Spectra of Fissionable Isotopes," in Proc. Conf "50 Years with Nuclear Fission," Washington, D.C., April 25-28, 1989, J. W. Behrens and A. D. Carlson, Eds.
- 10. G. Rudstam and T. R. England, "Test of Pre-ENDF/B-VI Decay and Fission Yields," Los Alamos National Laboratory report LA- -MS (to be published July 1990).




Fig. 1. <sup>235</sup>U independent yields (Los Alamos National Laboratory 1989).