CONF. 780622 -- 84

RECEIVED BY THO APR - 8 1980

LA-UR-78-427

TITLE: Reactivity Coefficients of Heavy Isotopes in LASL's Fast Critical Assemblies

AUTHOR(S): C. C. Byers

- G. E. Hansen
- J. J. Koelling E. A. Plassmann
- D. R. Smith

SUBMITTED TO: ANS Annual Meeting, San Diego, CA June 18-23, 1978

UBULATING in a series of the second of work schwards by access a filler (on periods) as a series of the terminal states to volument on any series (needed, top any of their omisingles, match any working, ended, or access or an even of any information apportance, product, or periods that the second of any information apportance, product, or periods they are series and any information apportance, product, or periods they are series and any information apportance, product, or periods they are series and any information apportance, product, or periods they are series and any information apportance, product, or periods they are series and any information apportance, periods and any series and any information apportance, periods and any series and apportance of the series (approximation of the series and apportance). The series of antibility series and apportance of antibility experimentation of series (approximation apportance) and any information of apportance of series (apportance). The series and apportance of automation apportance of antibility experimentation of series (apportance) and any approximation of apportance of apportance). The series of antibility experimentation of approximation apportance of antibility experimentation of apportance of a series of a new series of antibility experimentation of apportance of a series of a new series of apportance of a series of

- DISCLAIME

By acceptance of this article for publication, the publisher recognizes the Government's (license) rights in any copyright and the Government and its authorized representatives have unrestricted right to reproduce in whole or in part said article under any copyright secured by the publisher.

The Los Alumos Scientific Laboratory requests that the publisher identify this article as work performed under the nuspices of the USERDA.

alamos ntific laboratory SC of the University of California

LOS ALAMOS, NEW MEXICO 87545

An Allirmative Action/Equal Opportunity Employer

Form: No. 836 St. No. 2629 1/55

UNITED STATES ENERGY RESEARCH AND DEVELOPMENT ADMINISTRATION CONTRACT W-7405-ENG, 36

DISTRIBUTION OF THES POCUMENT IS UNLIMITED IC

REACTIVITY COEFFICIENTS OF HEAVY ISOTOPES IN LASL'S FAST CRITICAL ASSEMBLIES

by

C. C. Byers, G. E. Hansen, J. J. Koelling, E. A. Plassmann and D. R. Smith, University of California, Los Alamos Scientific Laboratory, Los Alamos, New Mexico 87545

We are attempting a systematic investigation of heavy isotope behavior in the Pajarito fast-neutron critical assemblies. Included will be central reactivity coefficient measurements and fission rate measurements in natural uranium reflected plutonium and 93%-enriched uranium metal assemblies and in Big Ten, a depleted uranium reflected 10%-enriched uranium metal assembly.

A few reactivity coefficients have recently been measured for ^{235}U , ^{237}Np , ^{238}Pu , ^{239}Pu , ^{242}Pu , and ^{241}Am . The experimental technique has been described by Stubbins, Barton, and Loandier,⁽¹⁾ and preliminary results are listed in the Table.

Similar data for some c? these isotopes in unreflected plutonium (Jezebel) and 93%-enriched uranium (Godiva) have previously been reported. (1-3)

Central reactivity coefficients are closely related to neutron net-production cross sections of the samples which in turn are related to criticality. By establishing the bias in the computed coefficients, one may obtain improved estimations of critical dimensions. Six group cross section sets, developed for the above isotopes, have been used in this fashion to give the following bare sphere critical mass estimates.

$$m_{c}^{(238}Pu) \approx 12 \text{ kg, } m_{c}^{(242}Pu) \approx 90 \text{ kg, and } m_{c}^{(241}Am) \approx 58 \text{ kg.}$$

It is hoped that these data will be useful to the standards work group which is currently developing a criticality safety standard for operations with the actinide elements.

TABLE

Central Reactivity Coefficient Ratios in Flattop-Pu, Flattop-U(93), and Big Ten

| Isotope | Flattop-Pu | Flattop-U(93.2) | Big Ten |
|-------------------|-------------|-----------------|-------------------|
| 235 _U | 0.524±0.007 | 0.528±0.010 | 0.585±0.006+.65% |
| 237 _{Np} | _ | 0.42 ±0.02 | 0.065±0.006 → 06- |
| 238 _{Pu} | 0.899±0.014 | 0.889±0.014 | 0.661±0.018 → 654 |
| 239 _{Pu} | ≡1.00 | ≡1.00 | ≡1.00 |
| 242 _{Pu} | 0.494±0.010 | 0.479±0.010 | - |
| 241 _{Am} | 0.605±0.011 | 0.530±0.010 | _ |

References

- W. F.Stubbins, D. M. Barton, and F. D. Lonadier, "The Neutron-Production Cross Section of ²³⁸Pu in a Fast Spectrum," Nucl. Sci. Eng. 25, 377-382, (1966).
- 2. D. M. Barton, "Central Reactivity Contributions of 244_{Cm}, ²³⁹Pu and ²³⁵U in a Bare Critical Assembly of Plutonium Metal," Nucl. Sci. Eng. 33, 51-55, (1968).

÷ ...

.

3. L. B. Engle, G. E. Hansen, and H. C. Paxton, "Reactivity Contributions of Various Materials in Topsy, Godiva, and Jezebel," Nucl. Sci. Eng. 8, 543-560, (1960).